

ON FACTORIALITY OF COX RINGS

IVAN V. ARZHANTSEV

ABSTRACT. Generalized Cox's construction associates with an algebraic variety a remarkable invariant – its total coordinate ring, or Cox ring. In this note we give a new proof of factoriality of the Cox ring when the divisor class group of the variety is finitely generated and free. The proof is based on a notion of graded factoriality. We show that if the divisor class group has torsion, then the Cox ring is again factorially graded, but factoriality may be lost.

1. INTRODUCTION

Let X be an irreducible normal algebraic variety over an algebraically closed field $\mathbb{K}$ with a free finitely generated divisor class group $\text{Cl}(X)$. Denote by $\text{WDiv}(X)$ the group of Weil divisors on X and fix a sublattice $K \subset \text{WDiv}(X)$ which maps onto $\text{Cl}(X)$ isomorphically. Following famous D. Cox's construction [4] from toric geometry, define the Cox ring of the variety X as

$$R(X) = \bigoplus_{D \in K} \mathcal{O}(X, D), \quad \text{where} \quad \mathcal{O}(X, D) = \{ f \in \mathbb{K}(X) \mid \text{div}(f) + D \geq 0 \}.$$

Multiplication on graded components of $R(X)$ coincides with multiplication in the field $\mathbb{K}(X)$ of rational functions, and extends to other elements by distributivity. It can be easily checked that the ring $R(X)$ depends on the choice of the lattice K only up to isomorphism (for a more general statement, see Proposition 3.2). An important property of $R(X)$ is that it is a factorial ring, see [2], [5]. Here we give a new proof of this result.

Theorem 1.1. *The ring $R(X)$ is factorial.*

Our main aim is to show that factoriality of $R(X)$ reflects the fact that any effective Weil divisor on X is a unique non-negative integral combination of prime divisors. This observation immediately implies that the multigraded ring $R(X)$ is "factorial on the set of homogeneous elements" (we call this a priori weaker property "graded factoriality"). Further, we prove that factoriality follows from graded factoriality. This approach is realized in Section 2.

In Section 3, the Cox ring $R(X)$ is defined in the case when the divisor class group has torsion. Following [2] and [7], we check that $R(X)$ is well-defined. Here the ring $R(X)$ is also factorially graded, but factoriality may be lost. The corresponding examples are given in Section 4, where we describe the Cox ring of a homogeneous space of an affine algebraic group.

The author is grateful to J. Hausen for valuable discussions which led to some of the results given below.

2000 *Mathematics Subject Classification.* 13F15, 14C20, 14L30.

The author gratefully acknowledges the support from INTAS YS 05-109-4958 and Deligne 2004 Balzan prize in mathematics.

2. PROOF OF THEOREM 1.1

We start with some elementary properties of multigraded algebras. Let R be a commutative associative algebra with unit over a field $\mathbb{K}$. Assume that R is graded by the lattice $\mathbb{Z}^n$,

$$R = \bigoplus_{u \in \mathbb{Z}^n} R_u.$$

Denote by $R^\times$ (resp. R^+) the multiplicative semigroup of invertible (resp. homogeneous) elements of R .

- Lemma 2.1.** (i) *Suppose that for any $a, b \in R^+$ the condition $ab = 0$ implies $a = 0$ or $b = 0$. Then R has no zero-divisors.*
(ii) *If R has no zero-divisors and for any $a, b \in R^+$ the condition $ab = 1$ implies $a, b \in R_0$, then $R^\times = R_0^\times$.*
(iii) *Suppose that R has no zero-divisors. If $a \in R^+$ and $a = bc$, then $b, c \in R^+$.*

Proof. Let us fix the lexicographic order on the lattice $\mathbb{Z}^n$. With any element $a \in R$ one associates two homogeneous elements $L(a)$ and $l(a)$, namely its leading and lowest terms. Clearly, $L(ab) = L(a)L(b)$ and $l(ab) = l(a)l(b)$. Now the statements of the lemma follow easily. $\square$

- Corollary 2.2.** (i) *The Cox ring $R(X)$ has no zero-divisors.*
(ii) *The semigroup $R(X)^\times$ coincides with $\mathcal{O}(X)^\times$, where $\mathcal{O}(X)$ is the algebra of regular functions on the variety X .*

Proof. Statement (i) follows from Lemma 2.1, (i), because $\mathbb{K}(X)$ has no zero-divisors. To prove (ii), note that if $f_1 \in \mathcal{O}(X, D_1)$, $f_2 \in \mathcal{O}(X, D_2)$ and $f_1 f_2 = 1$, then $D_1 + D_2 = 0$, $\text{div}(f_1) + \text{div}(f_2) = 0$, and the sum of the effective divisors $\text{div}(f_1) + D_1$ and $\text{div}(f_2) + D_2$ equals zero. Therefore $\text{div}(f_i) + D_i = 0$, $i = 1, 2$, and $D_i = 0$. Now one uses Lemma 2.1, (ii) and the equality $R(X)_0 = \mathcal{O}(X)$. $\square$

Definition 2.3. Let A be a finitely generated abelian group, and $R = \bigoplus_{u \in A} R_u$ be an A -graded algebra.

- A non-zero element $a \in R^+ \setminus R^\times$ is called *h-irreducible*, if the condition $a = bc$, $b, c \in R^+$ implies that either b or c is invertible.
- An A -graded algebra R is said to be *factorially graded*, if any its non-zero non-invertible homogeneous element may be expressed as a product of h-irreducible elements, and such an expression is unique up to association and renumbering.

Remark 2.4. Assume that $R = \bigoplus_{u \in \mathbb{Z}^n} R_u$ has no zero-divisors. It follows from Lemma 2.1 (iii) that if R is factorial, then it is factorially graded.

Proposition 2.5. *The Cox ring $R(X) = \bigoplus_{D \in K} \mathcal{O}(X, D)$ is factorially graded.*

Proof. Effective Weil divisors on X are in one-to-one correspondence with classes of associated elements of $R(X)^+$, and the product of homogeneous elements corresponds to the sum of divisors. This shows that classes of h-irreducible elements of the ring $R(X)$ correspond to prime divisors. Since any effective Weil divisor is a unique non-negative integral combination of prime divisors, the ring $R(X)$ is factorially graded. $\square$

Below we shall need the following well-known lemma. For convenience of the reader we give a short proof, cf. [11, Prop. 17.1].

Lemma 2.6. *Let T be an algebraic torus and Z a normal algebraic variety with a regular T -action. Then any Weil divisor on Z is linearly equivalent to a T -invariant Weil divisor.*

Proof. By normality, the singular locus of Z has codimension ≥ 2 , and one may assume that Z is smooth. Then any Weil divisor D on Z is Cartier. We may assume that D is effective. There exists a T -linearization of the corresponding line bundle, which defines a structure of a rational T -module on $\mathcal{O}(Z, D)$ [8, 2.4]. Any T -eigenvector in $\mathcal{O}(Z, D)$ represents a T -invariant divisor equivalent to D . $\square$

Again by normality, the passage from X to its smooth locus does not change the Cox ring. Further we shall assume that X is smooth. Following [2], let us introduce the "universal torsor" $\hat{X} \rightarrow X$ over X . Consider the K -graded sheaf of $\mathcal{O}_X$ -algebras

$$\mathcal{R}_X = \bigoplus_{D \in K} \mathcal{O}(D), \quad \text{where} \quad \mathcal{O}(U, D) = \{ f \in \mathbb{K}(X) \mid [\operatorname{div}(f) + D]|_U \geq 0 \},$$

and the relative spectrum $\hat{X} = \operatorname{Spec}_X(\mathcal{R}_X)$ of this sheaf over X . Clearly, $R(X) = H^0(X, \mathcal{R}_X) = \mathcal{O}(\hat{X})$. The K -grading on $\mathcal{R}_X$ defines a regular action of the torus $T = \operatorname{Spec}(\mathbb{K}[K])$ on $\hat{X}$, and the canonical affine morphism $p : \hat{X} \rightarrow X$ given by inclusion $\mathcal{O}_X \subset \mathcal{R}_X$ is T -invariant. Since all divisors on X are Cartier, p is a locally trivial fibration with T as a fiber. In particular, $\hat{X}$ is smooth. Fix an open affine covering $\{U_i\}$ of X . Each divisor $D_i = X \setminus U_i$ corresponds to an element $f_i \in R(X)^+$. The section f_i of the sheaf $\mathcal{R}_X$ is invertible exactly over U_i . Therefore $p^{-1}(U_i)$ coincides with $\hat{X}_{f_i} = \{z \in \hat{X} : f_i(z) \neq 0\}$. We get an open affine covering $\{\hat{X}_{f_i}\}$ of $\hat{X}$, where $f_i \in \mathcal{O}(\hat{X})$. It is easy to deduce from this that $\hat{X}$ is a quasi-affine variety, see. [6, Ch. 2, App., Lemma 8].

We are ready to finish with the proof of Theorem 1.1. Assume that the ring $R(X)$ is not factorial. Then $R(X)$ contains a non-principal prime ideal of height one. Since $R(X) = \mathcal{O}(\hat{X})$, it is a Krull ring, and its prime ideals of height one are in bijection with essential discrete valuations of the ring $R(X)$, see [10, I.3]. On the other hand, prime divisors on a normal quasi-affine variety are in bijection with essential discrete valuations of the ring of regular functions. Consequently, there is a non-principal divisor on $\hat{X}$. By Lemma 2.6, one may assume that there is a prime T -invariant non-principal divisor on $\hat{X}$. The corresponding ideal $I \triangleleft R(X)$ contains a homogeneous, and even an $\mathfrak{h}$ -irreducible element. Suppose that for some $a, b \in R(X)$ the product ab is divisible by p , but any homogeneous component of both a and b is not divisible by p . Since $R(X)$ is factorially graded, one considers the product of leading components of a and b and comes to contradiction. Thus the ideal $(p) \triangleleft R(X)$ is prime. The inclusion $(p) \subseteq I$ implies the equality $(p) = I$. This contradiction completes the proof of Theorem 1.1.

We finish this section with the following observation.

Proposition 2.7. *Let $R = \bigoplus_{u \in \mathbb{Z}^n} R_u$ be a multigraded finitely generated $\mathbb{K}$ -algebra without zero divisors. Assume that R is factorially graded. Then R is factorial.*

Proof. Let us show that R integrally closed. The multigrading defines actions of an n -dimensional torus T on the algebra R and its quotient field $\operatorname{Quot}(R)$ by automorphisms. It is known that the integral closure $\bar{R}$ of R in $\operatorname{Quot}(R)$ is a T -invariant subalgebra, and the T -action defines a structure of a rational T -module

on $\overline{R}$. In particular, $\overline{R}$ is $\mathbb{Z}^n$ -graded, and R is its homogeneous subalgebra. Any homogeneous element $r \in \overline{R}$ may be expressed as $r = \frac{r_1}{r_2}$, $r_1, r_2 \in R^+$. Since $R(X)$ is factorially graded and $\frac{r_1}{r_2}$ is integral over R , one has $\frac{r_1}{r_2} \in R^+$ and $\overline{R} = R$.

It is well known that R is factorial if and only if $\text{Cl}(Z) = 0$ for the normal affine variety $Z = \text{Spec}(R)$. The arguments given above show that any prime T -invariant divisor on Z is principal, and Lemma 2.6 completes the proof. $\square$

Remark 2.8. As in the proof of Proposition 2.7, one may show that in characteristic zero a factorially graded finitely generated algebra $R = \bigoplus_{u \in A} R_u$ without zero divisors is integrally closed for any finitely generated abelian group A .

3. TORSION IN THE DIVISOR CLASS GROUP

In this section we define the Cox ring $R(X)$ for a variety X with arbitrary finitely generated divisor class group and check that $R(X)$ is well-defined, compare [2], [7].

Let $S \subset \text{WDiv}(X)$ be a finitely generated subgroup that projects to $\text{Cl}(X)$ surjectively. Consider a ring

$$T_S(X) = \bigoplus_{D \in S} \mathcal{O}(X, D).$$

Let $S^0 \subset S$ be the kernel of the projection $S \rightarrow \text{Cl}(X)$. Take compatible bases $D_1, \dots, D_s$ in S and $D_1^0 = d_1 D_1, \dots, D_r^0 = d_r D_r$ in S^0 , $r \leq s$.

We call a family of rational functions $\mathcal{F} = \{F_D \in \mathbb{K}(X)^\times : D \in S^0\}$ *coherent*, if $\text{div}(F_D) = D$ and $F_{D+D'} = F_D F_{D'}$ (the term "shifting family" was used in [2] for a similar notion). Obviously, the family $\mathcal{F}$ is defined by $F_{D_i^0}$, $i = 1, \dots, r$: if $D = a_1 D_1^0 + \dots + a_r D_r^0$, then $F_D = F_{D_1^0}^{a_1} \dots F_{D_r^0}^{a_r}$. Let us fix a coherent family $\mathcal{F}$.

Let $D_1, D_2 \in S$ and $D_1 - D_2 \in S^0$. A map $f \mapsto F_{D_1 - D_2} f$ is an isomorphism between vector spaces $\mathcal{O}(X, D_1)$ and $\mathcal{O}(X, D_2)$. One easily checks, that the linear span of elements $f - F_{D_1 - D_2} f$ over all D_1, D_2 with $D_1 - D_2 \in S^0$ and all $f \in \mathcal{O}(X, D_1)$ is an ideal $I(S, \mathcal{F})$ of $T_S(X)$. Define the Cox ring of the variety X as

$$R_{S, \mathcal{F}}(X) = T_S(X) / I(S, \mathcal{F}).$$

Since $D/D^0 \cong \text{Cl}(X)$, the ring $R_{S, \mathcal{F}}(X)$ carries a natural $\text{Cl}(X)$ -grading.

Lemma 3.1. *Assume that $\mathcal{O}(X)^\times = \mathbb{K}^\times$. Then the ring $R_{S, \mathcal{F}}(X)$ does not depend on a choice of $\mathcal{F}$ up to isomorphism.*

Proof. The set of functions $F_{D_1^0}, \dots, F_{D_r^0}$ is defined up to transformations $F_{D_i^0} \rightarrow \gamma_i F_{D_i^0}$, $\gamma_i \in \mathbb{K}^\times$. Fix elements $\alpha_i \in \mathbb{K}^\times$, $i = 1, \dots, r$, such that $\alpha_i^{d_i} = \gamma_i$, and set $\alpha_{r+1} = \dots = \alpha_s = 1$. Then the desired isomorphism of the quotient-rings is induced by an automorphism $T_S(X) \rightarrow T_S(X)$, which acts on the component $\mathcal{O}(X, D)$, $D = a_1 D_1 + \dots + a_s D_s$, via multiplication by $\alpha_1^{a_1} \dots \alpha_s^{a_s}$. $\square$

Proposition 3.2. *Assume that $\mathcal{O}(X)^\times = \mathbb{K}^\times$. Then the ring $R_S(X)$ does not depend on a choice of S up to isomorphism.*

Proof. Let $M \subset \text{WDiv}(X)$ be another finitely generated subgroup that projects to $\text{Cl}(X)$ surjectively. One may assume that $\text{rk}(S) \geq \text{rk}(M)$.

Lemma 3.3. *There is a surjective homomorphism $\phi : S \rightarrow M$ such that the following diagram is commutative:*

$$\begin{array}{ccc} S & \xrightarrow{\phi} & M \\ & \searrow & \swarrow \\ & \text{Cl}(X) & \end{array}$$

Proof. Take compatible bases $D_1, \dots, D_s$ in S and $d_1 D_1, \dots, d_r D_r$ in S^0 , and also $M_1, \dots, M_k$ in M and $m_1 M_1, \dots, m_p M_p$ in M^0 , such that d_i (resp. m_i) is divisible by d_{i+1} (resp. by m_{i+1}). The condition $S/S^0 \cong \text{Cl}(X) \cong M/M^0$ implies that $s - r = k - p$, and the sets $(d_1, \dots, d_r)$ and $(m_1, \dots, m_p)$ may differ only in the final group of units. It remains to put $\phi(D_i) = M_i$ for $i = 1, \dots, p$, $\phi(D_i) = M_{i-r+p}$ for $i = r + 1, \dots, s$, and $\phi(D_i) = 0$ for other i . $\square$

We return to the proof of Proposition 3.2. Fix a coherent family $\mathcal{F}$ for the subgroup $(S + M)^0 \subset S + M$, and define a surjective homomorphism $\Phi : T_S(X) \rightarrow T_M(X)$ that sends a homogeneous component $\mathcal{O}(X, D)$ to $\mathcal{O}(X, \phi(D))$ as $\Phi(f) = F_{D-\phi(D)} f$. The kernel of Φ is contained in the ideal $I(S, \mathcal{F}|_{S^0})$, and the ideal $I(S, \mathcal{F}|_{S^0})$ itself maps surjectively to $I(M, \mathcal{F}|_{M^0})$. This shows that Φ defines a homomorphism $R_S(X) \rightarrow R_M(X)$ that, in fact, is an isomorphism. $\square$

Note that homogeneous elements of the ring $T_S(X)$ from different components corresponding to D_1 and D_2 with $D_1 - D_2 \in S^0$ may define the same effective divisor on X . However, after factorization by the ideal $I(S, \mathcal{F})$ effective Weil divisors on X are again in bijection with association classes in $R(X)^+$.

Proposition 3.4. *The ring $R(X)$ is factorially graded.*

Proof. The statement follows from the proof of Proposition 2.5. $\square$

Remark 3.5. By [5, Cor. 1.2], the ring $T_S(X)$ is factorial. As we shall see in the next section, this property may be lost for the ring $R(X)$ if the group $\text{Cl}(X)$ has torsion.

4. HOMOGENEOUS SPACES OF AN ALGEBRAIC GROUP

In this section we assume that the ground field $\mathbb{K}$ has characteristic zero. Let G be a connected affine algebraic group with $\text{Cl}(G) = 0$ and without nontrivial characters. Note that the first condition may be achieved by passing to a finite covering of a given group G [8, Prop. 4.6]. Denote by $\mathbb{X}(F)$ the group of characters of an algebraic group F . By Rosenlicht's Theorem, the condition $\mathbb{X}(G) = 0$ is equivalent to $\mathcal{O}(G)^\times = \mathbb{K}^\times$.

Let H be a closed subgroup of G . The homogeneous space G/H admits a canonical structure of a smooth quasi-projective algebraic G -variety. In [9], it was proved that $\text{Cl}(G/H) \cong \text{Pic}(G/H) \cong \mathbb{X}(H)$. Let us recall how to establish the last isomorphism.

Any character $\chi \in \mathbb{X}(H)$ defines a one-dimensional H -module $\mathbb{K}_\chi$. Consider a homogeneous fiber bundle

$$L_\chi = G \times_H \mathbb{K}_\chi := (G \times \mathbb{K}_\chi)/H, \quad h \cdot (g, a) := (gh^{-1}, \chi(h)a).$$

It is shown in [9] that the projection $L_\chi \rightarrow G/H$ is a G -linearized line bundle over G/H , and $L_{\chi_1} \otimes L_{\chi_2} \cong L_{\chi_1 + \chi_2}$. Moreover, the map $\chi \rightarrow L_\chi$ defines an isomorphism between $\mathbb{X}(H)$ and $\text{Pic}(G/H)$.

Since $\text{Cl}(G) = 0$, the pull-back of the line bundle L_χ with respect to the projection $G \rightarrow G/H$ is a trivial G -linearized line bundle on G . This allows to identify the space of sections $H^0(X, L_\chi)$ with the following subspace of $\mathcal{O}(G)$:

$$\mathcal{O}(G)_\chi^{(H)} := \{f \in \mathcal{O}(G) : f(gh^{-1}) = \chi(h)f(g) \text{ for all } h \in H, g \in G\}.$$

The tensor product of sections corresponds to the product in $\mathcal{O}(G)$.

Consider a subgroup $H_1 = \cap_{\chi \in \mathbb{X}(H)} \text{Ker}(\chi)$. The next theorem delivers an effective description of the Cox ring of the homogeneous space G/H , see also [1, Lemma 3.14].

Theorem 4.1. *Let G be a connected affine algebraic group with $\mathbb{X}(G) = 0$ and $\text{Cl}(X) = 0$, and H be a closed subgroup of G . Then*

$$R(G/H) \cong \mathcal{O}(G/H_1).$$

Proof. The diagonalizable group $Q = H/H_1$ acts on G/H_1 by right multiplication. Since $\mathbb{X}(H) = \mathbb{X}(Q)$ and any rational Q -module is a direct sum of one-dimensional submodules, we get $\mathcal{O}(G/H_1) = \bigoplus_{\chi \in \mathbb{X}(H)} \mathcal{O}(G)_\chi^{(H)}$.

Weil divisors on G/H are in bijection with lines generated by Q -semiinvariants $f \in \text{Quot}(\mathcal{O}(G/H_1)) \subseteq \mathbb{K}(G)$. Further, effective divisors correspond to semiinvariants $f \in \mathcal{O}(G/H_1)$. Let us choose a multiplicative finitely generated group of semiinvariants in $\text{Quot}(\mathcal{O}(G/H_1))$ whose weights run through the whole group $\mathbb{X}(Q)$. One may identify this subgroup with $S \subset \text{WDiv}(G/H)$ and check easily that the Cox ring associated with S is isomorphic to $\bigoplus_{\chi \in \mathbb{X}(H)} \mathcal{O}(G)_\chi^{(H)} \subset \mathcal{O}(G)$. $\square$

For a connected H , the group Q is a torus and $\mathbb{X}(H)$ is free. Moreover, in this case $\mathbb{X}(H_1) = 0$, thus $\text{Cl}(G/H_1) = 0$ and the ring $R(G/H)$ is factorial (it follows also from Theorem 1.1).

For a disconnected H , the character group $\mathbb{X}(H_1)$ may be non-trivial.

Example 4.2. Let $G = \text{SL}(2)$ and H be the normalizer N of a maximal torus $T \subset \text{SL}(2)$. Here $\mathbb{X}(H)$ is isomorphic to the cyclic group $\mathbb{Z}_2$ of order 2, $H_1 = T$, and $\mathbb{X}(H_1) \cong \mathbb{Z}$. This shows that the Cox ring $R(\text{SL}(2)/N) \cong \mathcal{O}(\text{SL}(2)/T)$ is not factorial. The space $\text{SL}(2)/N$ is a smooth affine surface X with $\text{Cl}(X) \cong \mathbb{Z}_2$, the ring $R(X)$ is isomorphic to

$$\mathbb{K}[x_1, x_2, x_3] / \langle x_2^2 - x_1x_3 - 1 \rangle,$$

and the $\mathbb{Z}_2$ -grading on $R(X)$ is given by $\deg(x_1) = \deg(x_2) = \deg(x_3) = \bar{1}$.

One may propose several ways to associate a factorial ring with the surface X . Firstly, consider a subgroup $S \subset \text{WDiv}(X)$ generated by a non-principal divisor. The ring $T_S(X)$ is isomorphic to

$$(\mathbb{K}[x_1, x_2, x_3, t, t^{-1}] / \langle x_2^2 - x_1x_3 - 1 \rangle)^{\mathbb{Z}_2},$$

where $\mathbb{Z}_2$ acts on the variables x_1, x_2, x_3, t, t^{-1} via multiplication by -1 .

Secondly, the space $\text{SL}(2)/N$ admits a wonderful (the term is due to D. Luna) $\text{SL}(2)$ -equivariant embedding in $\mathbb{P}^2$, and $R(\mathbb{P}^2) \cong \mathbb{K}[x_1, x_2, x_3]$. The Cox ring of a wonderful embedding of any spherical homogeneous space is described in [3].

Example 4.3. Using the construction of Example 4.2, one may find a smooth affine variety X with $\mathrm{Cl}(X) \cong A$ and a non-factorial Cox ring for any non-free finitely generated abelian group A . Indeed, let $A \cong \mathbb{Z}_{d_1} \oplus \cdots \oplus \mathbb{Z}_{d_s} \oplus \mathbb{Z}^n$. Put $X = G/H$, where G is a direct product of $s + n$ simple groups,

$$G = \mathrm{SL}(d_1) \times \cdots \times \mathrm{SL}(d_s) \times \mathrm{SL}(2) \times \cdots \times \mathrm{SL}(2),$$

and $H = H(1) \times \cdots \times H(s) \times T \times \cdots \times T$, where $H(i)$ is an extension of a maximal torus of the group $\mathrm{SL}(d_i)$ by elements of its normalizer that act as degrees of one d_i -cycle. In this case, $\mathbb{X}(H) \cong A$ and $\mathbb{X}(H_1) \cong \mathbb{Z}^{d_1 + \cdots + d_s - s}$.

REFERENCES

- [1] I.V. Arzhantsev, J. Hausen, On embeddings of homogeneous spaces with small boundary. *J. Algebra* **304:2** (2006), 950–988
- [2] F. Berchtold, J. Hausen, Homogeneous coordinates for algebraic varieties. *J. Algebra* **266:2** (2003), 636–670
- [3] M. Brion, The total coordinate ring of a wonderful variety. *J. Algebra* **313** (2007), 61–99
- [4] D.A. Cox, The homogeneous coordinate ring of a toric variety. *J. Alg. Geom.* **4** (1995), 17–50
- [5] J. Elizondo, K. Kurano, K. Watanabe, The total coordinate ring of a normal projective variety. *J. Algebra* **276** (2004), 625–637
- [6] F.D. Grosshans, *Algebraic Homogeneous Spaces and Invariant Theory*, LNM **1673**, Springer-Verlag Berlin, 1997
- [7] J. Hausen, Cox rings and combinatorics II, preprint [arXiv:0801.3995](https://arxiv.org/abs/0801.3995)
- [8] F. Knop, H. Kraft, D. Luna, Th. Vust, Local properties of algebraic group actions. In: *Algebraische Transformationsgruppen und Invariantentheorie*, DMV Seminar **13**, Birkhäuser, Basel (1989), 63–76
- [9] V.L. Popov, Picard group of homogeneous spaces of linear algebraic groups and one-dimensional homogeneous vector bundles. *Izvestia Acad. Nauk SSSR, Ser. Mat.* **38:2** (1974), 294–322; English transl.: *Math. USSR-Izv.* **8:2** (1974), 301–327
- [10] P. Samuel, *Lectures on unique factorization domains*, Tata Inst. Fund. Res., Bombay, 1964
- [11] D. A. Timashev, Homogeneous spaces and equivariant embeddings, preprint [math.AG/0602228](https://arxiv.org/abs/math/0602228)

DEPARTMENT OF HIGHER ALGEBRA, FACULTY OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY, LENINSKIE GORY, GSP-2, MOSCOW, 119992, RUSSIA

E-mail address: arjantse@mccme.ru